

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 June 2002 (27.06.2002)

PCT

(10) International Publication Number
WO 02/50100 A2

(51) International Patent Classification⁷: **C07K 5/00**

(21) International Application Number: **PCT/EP01/14821**

(22) International Filing Date:
17 December 2001 (17.12.2001)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
0031068.0 19 December 2000 (19.12.2000) **GB**

(71) Applicants (for all designated States except US):
SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
CAMBRIDGE UNIVERSITY TECHNICAL SERVICES LTD [GB/GB]; The Old Schools, Cambridge University (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CAMILLERI, Patrick** [MT/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). **KIRBY, Anthony, John** [GB/GB]; University Of Cambridge, Department Of Chemistry, Chemical Laboratory, Lensfield Road, Cambridge, CB2 1EW (GB). **PERRIN, Christele** [FR/GB]; University of Cambridge, Department of Chemistry, Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW (GB). **RON SIN, Gaël, Alain, Bertrand** [FR/GB]; University of Cambridge, Department

of Chemistry, Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW (GB). **GUEDAT, Philippe** [FR/FR]; LIPHA S.A., Centre de Recherche et Developpement de Lyon Lacassagne, 115, avenue Lacassagne, F-69424 Lyon Cedex 03 (FR).

(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline, Corporate Intellectual Property (CN9.25.1), 980 Great West Road, Brentford, Middlesex TW8 9GS (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/50100 A2

(54) Title: **NOVEL COMPOUNDS**

(57) Abstract: Diaminodicarboxylic acid:peptide gemini surfactant compounds are disclosed. Uses of the diaminodicarboxylic acid: peptide-based gemini surfactant compounds and methods for their production are also disclosed.

BEST AVAILABLE COPY

Novel compounds

This invention relates to newly identified diaminodicarboxylic acid:peptide-based gemini surfactant compounds, to the use of such compounds and to their production. The invention also relates to the use of diaminodicarboxylic acid:peptide-based gemini compounds to facilitate the transfer of compounds into cells for drug delivery.

Surfactants are substances that markedly affect the surface properties of a liquid, even at low concentrations. For example surfactants will significantly reduce surface tension when dissolved in water or aqueous solutions and will reduce interfacial tension between two liquids or a liquid and a solid. This property of surfactant molecules has been widely exploited in industry, particularly in the detergent and oil industries. In the 1970s a new class of surfactant molecule was reported, characterised by two hydrophobic chains with polar heads which are linked by a hydrophobic bridge (Deinega, Y *et al.*, *Kolloidn. Zh.* 36, 649, 1974). These molecules, which have been termed "gemini" (Menger, FM and Littau, CA, *J. Am. Chem. Soc.* 113, 1451, 1991), have very desirable properties over their monomeric equivalents. For example they are highly effective in reducing interfacial tension between oil and water based liquids and have a very low critical micelle concentration.

Cationic surfactants have been used *inter alia* for the transfection of polynucleotides into cells in culture, and there are examples of such agents available commercially to scientists involved in genetic technologies (for example the reagent TfxTM-50 for the transfection of eukaryotic cells available from Promega Corp. WI, USA).


The efficient delivery of DNA to cells *in vivo*, either for gene therapy or for antisense therapy, has been a major goal for some years. Much attention has concentrated on the use of viruses as delivery vehicles, for example adenoviruses for epithelial cells in the respiratory tract with a view to corrective gene therapy for cystic fibrosis (CF). However, despite some evidence of successful gene transfer in CF patients, the adenovirus route remains problematic due to inflammatory side-effects and limited transient expression of the transferred gene. Several alternative methods for *in vivo* gene delivery have been investigated, including studies using cationic surfactants. Gao, X *et al.* (1995) *Gene Ther.* 2, 710-722 demonstrated the feasibility of this approach with a normal human gene for CF transmembrane conductance regulator (CFTR) into the respiratory epithelium of CF mice using amine carrying cationic lipids. This group followed up with a liposomal CF gene therapy trial which, although only partially successful, demonstrated the potential for this

approach in humans (Caplen, NJ. *et al.*, *Nature Medicine*, 1, 39-46, 1995). More recently other groups have investigated the potential of other cationic lipids for gene delivery, for example cholesterol derivatives (Oudrhiri, N *et al.* *Proc. Natl. Acad. Sci.* 94, 1651-1656, 1997). This limited study demonstrated the ability of these cholesterol based compounds to facilitate the transfer of genes into epithelial cells both *in vitro* and *in vivo*, thereby lending support to the validity of this general approach.

These studies, and others, show that in this new field of research there is a continuing need to develop novel low-toxicity surfactant molecules to facilitate the effective transfer of polynucleotides into cells both *in vitro* for transfection in cell-based experimentation and *in vivo* for gene therapy and antisense treatments. The present invention seeks to overcome the difficulties exhibited by existing compounds.

Recently a number of peptide-based gemini surfactants having gene transfection properties were disclosed in WO99/29712 (SmithKline Beecham).

The invention relates to diaminodicarboxylic acid:peptide-based gemini compounds having a diaminodicarboxylic acid backbone and conforming to the general structure of formula (I):



20

(I)

where X = (CH₂)_{n2}, n₂ is 1 to 8 and n₁ is 0; or

where X = NHC(O)(CH₂)_{n3}C(O)NH, n₃ is 1 to 8 and n₁ is 2 to 4; or

25

where X = (CH₂)_{n4}NHC(O)(CH₂)_{n5}C(O)NH(CH₂)_{n4}, n₄ is 2 to 4, n₅ is 1 to 8 and n₁ is 0;

and

where R₃, R₄, R₅ and R₆ is hydrogen;

5 or

where R₃ and R₅ is hydrogen and R₄ and R₆ which may be the same or different are peptide groups formed from one or more amino acids linked together by amide (CONH) bonds and further linked to the diaminodicarboxylic acid backbone by amide bonds, in a linear or branched manner, having the general formula (II):

10



15 where the values for p₁ and p₂, which may be the same or different, are from 0 to 5, preferably 1; and

the values for p₃ and p₄, which may be the same or different, are from 0 to 5, preferably 0; A₁, A₃ and A₄, which may be the same or different, is an amino acid selected from serine, lysine, ornithine, threonine, histidine, cysteine, arginine and tyrosine; and

20 A₂ is an amino acid selected from lysine, ornithine and histidine;

or

where R₄ and R₆, which may be the same or different, is a group having the formula (III):

25



and R₃ and R₅, which may be the same or different, is a group having the formula (IV):

30



where p₁, p₂, p₃, and p₄ which may be the same or different, are 0 to 5;

and where m is 0 to 5; q is 1 to 5;

and where A1 to A4 are as defined above;

and R₁ and R₂ are saturated or unsaturated aminohydrocarbyl groups having up to 32 carbon
5 atoms and linked to the diaminodicarboxylic acid backbone by an amide bond;


or

a salt, preferably a pharmaceutically acceptable salt thereof.


Preferably, the compound is symmetrical, that is R₁ and R₂ are the same, R₃ and
10 R₅ are the same, and R₄ and R₆ are the same.

In a preferred embodiment A1 is serine or threonine, preferably serine. Preferably
A3 and A4 are lysine, ornithine, histidine or arginine.

In a further preferred embodiment the aminohydrocarbyl group is selected from:



15



Compounds of the present invention may be prepared from readily available starting materials using synthetic peptide chemistry well known to the skilled person. The scheme shown in Figure 1 shows a general scheme for the synthesis of the compounds of the invention wherein the aminohydrocarbyl groups are linked to the diaminodicarboxylic acid moiety by amide bonds, the scheme shown in Figure 2 shows a general scheme for the synthesis of the compounds of the invention wherein the head group is a diacid linked to the α -amino group of diaminoacid moiety by amide bonds and the scheme shown in Figure 3 shows a general scheme for the synthesis of the compounds of the invention wherein the head group is a diacid linked to the non- α -amino group of a diaminoacid moiety by amide bonds.

Another aspect of the invention relates to methods for using diaminodicarboxylic acid:peptide-based gemini compounds. Such uses include facilitating the transfer of oligonucleotides and polynucleotides into cells for antisense, gene therapy and genetic immunisation (for the generation of antibodies) in whole organisms. Other uses include employing the compounds of the invention to facilitate the transfection of polynucleotides into cells in culture when such transfer is required, in, for example, gene expression studies and antisense control experiments among others. The polynucleotides can be mixed with the compounds, added to the cells and incubated to allow polynucleotide uptake. After further incubation the cells can be assayed for the phenotypic trait afforded by the transfected DNA, or the levels of mRNA expressed from said DNA can be determined by Northern blotting or by using PCR-based quantitation methods for example the Taqman[®] method (Perkin Elmer,

Connecticut, USA). Compounds of the invention offer a significant improvement, typically between 3 and 6 fold, in the efficiency of cellular uptake of DNA in cells in culture, compared with compounds in the previous art. In the transfection protocol, the gemini compound may be used in combination with one or more supplements to increase the efficiency of transfection. Such supplements may be selected from, for example:

- (i) a neutral carrier, for example dioleoyl phosphatidylethanolamine (DOPE) (Farhood, H., *et al* (1985) *Biochim. Biophys. Acta* 1235 289);
- (ii) a complexing reagent, for example the commercially available PLUS reagent (Life Technologies Inc. Maryland, USA) or peptides, such as polylysine or polyornithine peptides or peptides comprising primarily, but not exclusively, basic amino acids such as lysine, ornithine and/or arginine. The list above is not intended to be exhaustive and other supplements that increase the efficiency of transfection are taken to fall within the scope of the invention.

In still another aspect, the invention relates to the transfer of genetic material in gene therapy using the compounds of the invention.

Yet another aspect of the invention relates to methods to effect the delivery of non-nucleotide based drug compounds into cells *in vitro* and *in vivo* using the compounds of the invention.

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Amino acid" refers to dipolar ions (zwitterions) of the form $^+H_3NCH(R)CO_2^-$. They are differentiated by the nature of the group R, and when R is different from hydrogen can also be asymmetric, forming D and L families. There are 20 naturally occurring amino acids where the R group can be, for example, non-polar (e.g. alanine, leucine, phenylalanine) or polar (e.g. glutamic acid, histidine, arginine and lysine). In the case of un-natural amino acids R can be any other group which is not found in the amino acids found in nature.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically,

double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications have been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.


"Transfection" refers to the introduction of polynucleotides into cells in culture using methods involving the modification of the cell membrane either by chemical or physical means. Such methods are described in, for example, Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). The polynucleotides may be linear or circular, single-stranded or double-stranded and may include elements controlling replication of the polynucleotide or expression of homologous or heterologous genes which may comprise part of the polynucleotide.

The invention will now be described by way of the following examples.

EXAMPLES

Example 1

RG 00/184




25

D,L- α,ϵ -diaminopimelic acid (29.0 mmol ; 5.52 g) was dissolved in THF / water 1/1 (50 ml) then NaOH (11.6 mmol; 2.55 g ; 2.2 eq.) and Boc_2O (11.6 mmol; 13.94 g ; 2.2 eq.) were added. The mixture was stirred overnight at room temperature. Most of the THF

was removed and the mixture was acidified to pH 2 with HCl 3 M. The precipitate was extracted twice with chloroform then organic layers were combined and washed successively with water and brine, dried over anhydrous sodium sulfate and evaporated to yield the bis-protected compound (9.52 g, 84 %).


5

Example 2**RG 00/190**

- 10 The bis-Bocdiaminopimelic acid (7.07 g, 18.1 mmol) of example 1 was dissolved in THF (160 mL) then N-hydroxysuccinimide (4.34 g, 37.7 mmol, 2.1 eq.) and DCC (7.62 g, 36.9 mmol, 2.04 eq.) were added. The mixture was stirred 20 h at room temperature. The precipitate was filtered off and washed with EtOAc. Solvents were removed and the residue redissolved in EtOAc, cooled to 0°C and filtered. After evaporation of the
- 15 solvent, Et₂O is added to precipitate the solid. After filtration of Et₂O, a white solid (9.81 g, 93 %) is obtained. Mass spectrum (+ESI): 607.2231 (M+Na).

Example 3**RG00/219**

20




The bis-activated diaminopimelic derivative (1.48g, 2.53 mmol) of example 2 was

dissolved in THF (60 mL) then oleylamine 2.2 eq. (1.49 g, 5.57 mmol) and K_2CO_3 2.2 eq. (0.77 g, 5.57 mmol) in 5 ml of water were added. The mixture was left stirring at room temperature for 24 h. Most of the THF was removed by evaporation then water (100 mL) was added. The mixture was extracted with $CHCl_3$ (2 x 60 ml). The combined
5 organic layers were washed with brine, dried over Na_2SO_4 and evaporated. The residue was purified by chromatography on silica gel with ether / dichloromethane (1/2, v/v, rf: 0.3) to yield a white solid foam (1.68 g, 75 %). Mass spectrum (+ESI): 911.7554 (M+Na).

10 Example 4


RG 00/220



The di-Boc protected RG 00/219 of example 3 (1.50 g, 1.69 mmol) was stirred in a 1/1
15 mixture of CH_2Cl_2 /TFA (20 mL) for 1h. The solvent were evaporated. The oily residue was diluted in $CHCl_3$ and washed successively with 1M K_2CO_3 , water and brine, dried on sodium sulfate, filtered and evaporated to give a yellowish solid. Trituration in Et_2O followed by filtration gave a white powder (1.15 g, 99 %).

20 Example 5

PG991005.1




The diamine PG32788 100 mg (0.11 mmol) was dissolved in THF (10 ml) then potassium carbonate 2.2 eq. (34 mg; 0.24 mmol) in water (2 ml) and the *N* α ,*N* ϵ -bis-*tert*-butyl-carbamate-L-lysine-L-serine-*O*-succinimide (was built by the usual peptide synthesis) 2.05 eq. (0.22 mmol; 119 mg) in THF (8 ml) were added. The mixture was stirred overnight at room temperature. Most of the THF was removed then water (15 ml) was added and the precipitate was extracted with chloroform (2 x 25 ml). Organic phases were combined and washed with 4% NaHCO₃ (15 mL), water (15 mL), 4% citric acid (15 mL), water (15 mL), brine (15 mL), dried over sodium sulfate and concentrated to yield

10 160 mg (96 %) of coupling compound.

Example 6

GSC61




PG991005.1 from example 5 (155 mg, 0.102 mmol) was dissolved in a mixture of methanol / conc. HCl 1/1 (20 ml). The mixture was stirred 2 h at room temperature.

- 5 Solvent was removed and crude product was redissolved in water (80 ml), filtrated on sintered frit funnel (N° 3) then freeze dry to yield 108 mg (83 %) of GSC61.

Example 7

RG 00/265




10

- To a solution of adipoyl chloride (3.66g, 20 mmol) in tetrahydrofuran (180 ml) were added N-hydroxysuccinimide (4.60 g, 40 mmol, 2 eq.) and triethylamine (5.69 mL, 40 mmol, 2 eq.). After 24 h at room temperature, the solvent was evaporated and the residue was partitioned between dilute aqueous HCl and chloroform. The organic layer was
 15 extracted and washed successively with water and brine, dried over sodium sulphate, filtered and evaporated to give a white solid (5.99 g, 88 %).

NMR ^1H (CDCl_3): δ 1.69 (m, 2H, CH_2), 2.69 (m, 2H), 2.78 (s, 4H, succinimide)

Example 8


RG 00/274



- 5 To a stirred solution of adipoyl disuccinimidate (1.50 g, 4.41 mmol) in tetrahydrofuran (150 mL) were added the α - or ϵ -Boc-protected lysine (2.17 g, 8.82 mmol, 2 eq.) and potassium carbonate (1.30 g, 9.40 mmol, 2.1 eq.) in 30 ml of water. The reaction is stirred at room temperature for 20 h. Most of the THF was removed and the mixture was acidified to pH 2 with 3 M HCl. The precipitate was extracted twice with chloroform then
- 10 organic layers were combined and washed successively with water and brine, dried over anhydrous sodium sulfate and evaporated to give a white powder (2.66 g, 77 %).


Example 9

RG 00/285




To a solution of di-N-(N- ϵ -*tert*-butylcarbonate-lysiny)adipate (2.30 g, 3.82 mmol) in
5 tetrahydrofuran (180 mL) were added N-hydroxysuccinimide (0.90 g, 7.80 mmol, 2.05
eq.) and DCC (1.57 g, 7.66 mmol, 2 eq.). The mixture was stirred for 24 h at room
temperature and DCU was filtered and washed with EtOAc (3 x 30 mL). The solvents
were evaporated and the residue dissolved in EtOAc (40 mL) and the precipitate filtered.
After evaporation, the oily was cristallised in Et₂O to give a white powder (2.80 g, 92 %).

10

Example 10**RG 00/292**


To a solution of the succinimidyl ester RG 00/285 of example 9 (1.50 g, 1.88 mmol) in
5 THF (80 mL) were added oleylamine (1.01 g, 3.78 mmol, 2.02 eq.) and potassium
carbonate (0.53 g, 3.83 mmol, 2.1 eq.) in water (10 mL). The reaction was stirred for 20 h
at room temperature. Most of the THF was removed and the mixture was partitioned
between water and chloroform. The organic layer was extracted and washed successively
with water, 1M HCl, water and brine. After drying over sodium sulfate, filtration and
10 evaporation, the residue was purified by chromatography on silica gel in CHCl₃ / MeOH
(97/3, v/v, R_f : 0.33) to give an oil (1.35 g, 65 %). Mass spectrum (+ESI): 1123.89793
(M+Na).

Example 11**RG 00/296**

5 The di-Boc protected RG 00/292 of example 10 (1.20 g, 1.09 mmol) was stirred in a 1/1 mixture of CH₂Cl₂/TFA (20 mL) for 1h. The solvent were evaporated. The oily residue was diluted in CHCl₃ and washed successively with 1M K₂CO₃, water and brine, dried on sodium sulfate, filtered and evaporated to give a yellowish solid. Trituration in Et₂O followed by filtration gave a white powder (0.95 g, 97 %). Mass spectrum (+ESI): 901.82280 (MH⁺).

10


Example 12**RG 00/299**



To a solution of the diamino RG 00/296 from example 11 (228 mg, 0.25 mmol) and K_2CO_3 (74 mg, 0.54 mmol, 2.1 eq.) in a 9/1 THF/water mixture (40 mL, v/v) was added the activated peptide $Boc_2K-\epsilon-K(Boc)-\epsilon-K(Boc)-S-OSu$ (500 mg, 0.5 mmol, 2 eq.). The reaction was stirred for 24 h at room temperature. Most of the THF was removed under vacuum. A 4% Na_2CO_3 solution (10 mL) was added and the aqueous layer was extracted with $CHCl_3$ (3 x 40 mL). The combined organic layers were washed successively with water (20 mL), 4% citric acid (20 mL), water (20 mL), brine (20 mL), dried over sodium sulfate, filtered and evaporated to give a pale yellowish solid (543 mg, 81%). Mass spectrum (+ES): 2666.848 (M+Na).

Example 13

GSC 144




To a solution of Boc-protected gemini RG 00/299 from example 12 (500 mg, 0.19 mmol) in MeOH (10 mL) was added concentrated aqueous HCl (10 mL). The reaction
 5 was stirred for 1.5 h at room temperature. The solvents were evaporated and the residue dissolved in water (80 mL), filtered on a N°3 frit sinter. The aqueous layer was evaporated to dryness. The residue was dissolved in MeOH (4 mL) and Et₂O was added. The precipitate was collected to give a pale pink powder (348 mg, 86%). Mass spectrum (+ESI): 1844.446 (MH⁺).

10

Example 14


RG 00/348



To a solution of diamino derivative from example 13 (255 mg, 0.28 mmol) and potassium carbonate (82 mg, 0.59 mmol, 2.1 eq.) in a 9/1 mixture of THF/water (80 mL) was added *N*,*N*,*N*,*N*-bis-*tert*-butyl-carbamate-L-lysine-L-serine-*O*-succinimide (300 mg, 5.65 mmol, 2.05 eq.). The reaction was stirred at room temperature for 20 h. Most of the THF was removed by evaporation and the aqueous residue diluted with 4% NaHCO₃ (20 mL). The aqueous layer was then extracted twice with CHCl₃ (25 mL). The combined organic layers were washed successively with water (10 mL), 0.01 M HCl (10 mL), water (15 mL) and brine (20 mL), dried over sodium sulfate, filtered and evaporated to give a white solid (462 mg, 94%).

Example 15


GSC 150



To a solution of protected gemini RG 00/348 from example 14 (441 mg, 0.25 mmol) in MeOH (10 mL) was added concentrated HCl (10 mL). The mixture is then stirred for 1 h at room temperature. The solvents were evaporated to dryness. The residue was redissolved in water (80 mL) and filtered on a sintered frit funnel (N° 3) and evaporated using ethanol as co-solvent. The residue was then dissolved in MeOH (5 mL) and Et₂O was added until complete precipitation. The solid was isolated by decantation and dried under high vacuum to give a yellowish powder (341 mg, 91 %).

10 Example 16 – Compound GSC 112

Compound GSC 112 is an aminopimelic gemini compound, synthesised according to the schemes described herein and has the structure:



Example 17 - Transfection of recombinant plasmid expressing luciferase into cells using aminopimelic and adipate-Lysine:peptide-based gemini surfactant

5 compounds.

Transfection studies were performed using the adherent cell line CHO-K1 (Puck et al. 1958). Complete medium consisted of MEM alpha medium supplemented with 10 % v/v foetal bovine serum and 1x L-Glutamine. All media and supplements were obtained from Life Technologies.

10

***In Vitro* Gene Transfection.**

Cells were seeded into 96-well MTP plates (Nunc) 16-18 hours prior to transfection at an approximate density of 1×10^4 cells per well. For transfection, 0.064 μg of the luciferase reporter gene plasmid, pGL3-Control Vector (Promega) per well, was incubated with various concentrations of the gemini compounds GSC112 or GSC150 and complexing agents in a final volume of 100 μl . After 30 minutes incubation at RT, OPTI-MEM[®] medium (Life Technologies) was added to the transfection mixture and the solution placed on the cells (final volume per well: 100 μl). Following a 3 hour or over night incubation at 37°C, the transfection solution was replaced with complete medium and the cells incubated further at 37°C. Reporter gene assays were performed according to the manufacturer's guidelines (Roche Diagnostics) approximately 48 hours post transfection. Luminescence was measured in a Packard TopCount NXT Microplate Scintillation and Luminescence Counter.

20

Brief description of the drawings

Figure 1. General scheme for synthesis of compounds of the invention wherein the aminohydrocarbonyl groups are linked to the diaminodicarboxylic acid moiety by amide bonds.

Figure 2. General scheme for synthesis of compounds of the invention wherein the head group is a diacid linked to the α -amino group of the diaminoacid moiety by amide bonds.

Figure 3. General scheme for synthesis of compounds of the invention wherein the head group is a diacid linked to the non- α -amino group of a diaminoacid moiety by amide bonds.


Figure 4. Transfection of CHO-K1 cells with gemini surfactant GSC112. The numbers along the x axis refer to concentration of gemini compounds in μM . The block of 5 bars at the right of the chart show the data obtained when the DNA was premixed with polylysine. The block of 5 bars at the left side of the chart show data when no polylysine was used. The figures on the Y axis represent CPS (count per second) from the luciferase assay. The bars represent the mean CPS (count per second) of 4 experiments \pm the standard error of the mean.

Figure 5. Transfection of CHO-K1 cells with gemini surfactant GSC150. The numbers along the x axis refer to concentration of gemini compounds in μM . The block of 5 bars at the right of the chart show the data obtained when the DNA was premixed with polylysine. The block of 5 bars at the left side of the chart show data when no polylysine was used. The figures on the Y axis represent CPS (count per second) from the luciferase assay. The bars represent the mean CPS (count per second) of 4 experiments \pm the standard error of the mean.

CLAIMS

1. A diaminodicarboxylic acid:peptide-based gemini compound having a diaminodicarboxylic acid backbone and conforming to the general structure of formula (I):

5



(I)

10 where $\text{X} = (\text{CH}_2)_{n2}$, $n2$ is 1 to 8 and $n1$ is 0; or

where $\text{X} = \text{NHC}(\text{O})(\text{CH}_2)_{n3}\text{C}(\text{O})\text{NH}$, $n3$ is 1 to 8 and $n1$ is 2 to 4; or

15 where $\text{X} = (\text{CH}_2)_{n4}\text{NHC}(\text{O})(\text{CH}_2)_{n5}\text{C}(\text{O})\text{NH}(\text{CH}_2)_{n4}$, $n4$ is 2 to 4, $n5$ is 1 to 8 and $n1$ is 0;


and

where R_3 , R_4 , R_5 and R_6 is hydrogen;

20 or

where R_3 and R_5 is hydrogen and R_4 and R_6 which may be the same or different are peptide groups formed from one or more amino acids linked together by amide (CONH) bonds and further linked to the diaminodicarboxylic acid backbone by amide bonds, in a linear or branched manner, having the general formula (II):

25





where the values for p1 and p2, which may be the same or different, are from 0 to 5, preferably 1; and

- 5 the values for p3 and p4, which may be the same or different, are from 0 to 5, preferably 0;
 A1, A3 and A4, which may be the same or different, is an amino acid selected from serine, lysine, ornithine, threonine, histidine, cysteine, arginine and tyrosine; and
 A2 is an amino acid selected from lysine, ornithine and histidine;

or

- 10 where R4 and R6, which may be the same or different, is a group having the formula (III):



- 15 and R3 and R5, which may be the same or different, is a group having the formula (IV):




- 20 where p1, p2, p3, and p4 which may be the same or different, are 0 to 5;
 and where m is 0 to 5; q is 1 to 5;
 and where A1 to A4 are as defined above;

- and R1 and R2 are saturated or unsaturated aminohydrocarbyl groups having up to 32 carbon
 25 atoms and linked to the diaminodicarboxylic acid backbone by an amide bond;


or

a salt, preferably a pharmaceutically acceptable salt thereof.

2. A diaminodicarboxylic acid:peptide-based gemini compound according to claim 1 which is symmetrical, that is R_1 and R_6 are the same, R_2 and R_4 are the same, and R_3 and R_5 are the same.
- 5 3. A diaminodicarboxylic acid:peptide-based gemini compound according to claim 1 or 2 wherein in the peptide group of formula (II) p_1 and p_2 are both 1 and p_3 and p_4 are both 0.
4. A diaminodicarboxylic acid:peptide-based gemini compound according to any one
10 of claims 1 to 3 wherein the A_1 is serine.
5. A diaminodicarboxylic acid:peptide-based gemini compound according to any one of claims 1 to 4 wherein the A_2 is lysine.
- 15 6. A diaminodicarboxylic acid:peptide-based gemini compound according to claim 1 wherein the aminohydrocarbyl group is selected from:
- NH(CH₂)₁₁CH₃
 - NH(CH₂)₁₃CH₃
 - NH(CH₂)₁₅CH₃
 - NH(CH₂)₁₇CH₃
 - NH(CH₂)₁₉CH₃
 - NH(CH₂)₂₃CH₃
 - NH(CH₂)₈CH=CH(CH₂)₅CH₃
 - NH(CH₂)₈CH=CH(CH₂)₇CH₃
 - NH(CH₂)₈CH=CHCH₂CH=CH(CH₂)₄CH₃
 - NH(CH₂)₈(CH=CHCH₂)₃CH₃
 - NH(CH₂)₄CH=CH(CH₂CH=CH)₃(CH₂)₄CH₃
 - NH(CH₂)₈CH=CH(CH₂)₅CH₃ Trans
 - NH(CH₂)₈CH=CH(CH₂)₇CH₃ Trans
 - NH(CH₂)₉CHCH₃(CH₂)₇CH₃
7. A diaminodicarboxylic acid:peptide-based gemini compound according to claim 1
20 wherein the aminohydrocarbyl group is selected from:




8. The compound:




5

9. The compound:




10. The compound:




5

11. The compound GSC61 of formula:




12. The compound GSC 144 of formula:



5

13. The compound GSC150 of formula:



14. The use of a diaminodicarboxylic acid:peptide-based gemini compound according
5 to claim 1 to effect the delivery of polynucleotides into cells *in vitro* and *in vivo*.
15. The use of a diaminodicarboxylic acid:peptide-based gemini compound according
to claim 1 to effect the delivery of non-nucleotide based drug compounds into cells *in vitro*
and *in vivo*.
- 10 16. The use of a diaminodicarboxylic acid:peptide-based gemini compound according
to claim 14 or 15 wherein the compound is used in combination with one or more
supplements selected from the group consisting of:
- (i) a neutral carrier; or
- 15 (ii) a complexing reagent.
17. The use according to claim 16 wherein the neutral carrier is dioleoyl
phosphatidylethanolamine (DOPE).

18. The use according to claim 17 wherein the complexing reagent is selected from the group consisting of:
- i) PLUS reagent;
 - ii) a peptide comprising mainly basic amino acids;
 - 5 iii) a peptide consisting of basic amino acids;
 - iv) a peptide consisting of basic amino acids selected from lysine and arginine.

Figure 1




Figure 2




Figure 3






Figure 4

Figure 5



(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 June 2002 (27.06.2002)

PCT

(10) International Publication Number
WO 02/050100 A3

(51) International Patent Classification⁷: **C07K 5/02,**
C07C 237/00, 237/52

Lyon Lacassagne, 115, avenue Lacassagne, F-69424 Lyon
Cedex 03 (FR).

(21) International Application Number: **PCT/EP01/14821**

(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline, Corporate Intellectual Property (CN9.25.1), 980 Great West Road, Brentford, Middlesex TW8 9GS (GB).

(22) International Filing Date:
17 December 2001 (17.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0031068.0 19 December 2000 (19.12.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicants (*for all designated States except US*):
SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
CAMBRIDGE UNIVERSITY TECHNICAL SERVICES LTD [GB/GB]; The Old Schools, Cambridge University (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CAMILLERI, Patrick** [MT/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). **KIRBY, Anthony, John** [GB/GB]; University Of Cambridge, Department Of Chemistry, Chemical Laboratory, Lensfield Road, Cambridge, CB2 1EW (GB). **PERRIN, Christele** [FR/GB]; University of Cambridge, Department of Chemistry, Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW (GB). **RON SIN, Gaël, Alain, Bertrand** [FR/GB]; University of Cambridge, Department of Chemistry, Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW (GB). **GUEDAT, Philippe** [FR/FR]; LIPHA S.A., Centre de Recherche et Developpement de

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
12 September 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIAMINODICARBOXYLIC ACID: PEPTIDE-BASED GEMINI SURFACTANT COMPOUNDS USED FOR DRUG DELIVERY

(57) Abstract: Diaminodicarboxylic acid:peptide gemini surfactant compounds are disclosed. Uses of the diaminodicarboxylic acid: peptide-based gemini surfactant compounds and methods for their production are also disclosed.



WO 02/050100 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/14821

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K5/02 C07C237/00 C07C237/52

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 29712 A (CAMILLERI PATRICK ;RICE SIMON QUENTYN JOHN (GB); KREMER ANDREAS (G) 17 June 1999 (1999-06-17) cited in the application claims 1,21-31	1-7, 11-18
A	EP 0 394 111 A (CENTRE NAT RECH SCIENT) 24 October 1990 (1990-10-24) the whole document	1-18



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

19 June 2002

Date of mailing of the international search report

02/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

G. Willière

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/14821

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JENNINGS K ET AL: "The synthesis and aggregation properties of novel anionic gemini surfactant" CHEMICAL COMMUNICATIONS, ROYAL SOCIETY OF CHEMISTRY, GB, no. 18, 1 September 1998 (1998-09-01), pages 1951-1952, XP002096476 ISSN: 1359-7345 the whole document	1-18
A	WADHWA M S ET AL: "PEPTIDE-MEDIATED GENE DELIVERY: INFLUENCE OF PEPTIDE STRUCTURE ON GENE EXPRESSION" BIOCONJUGATE CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 8, no. 1, 1997, pages 81-88, XP000642731 ISSN: 1043-1802 the whole document	1-18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 01/14821

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9929712	A	17-06-1999	CA 2312941 A1	17-06-1999
			EP 1037903 A1	27-09-2000
			WO 9929712 A1	17-06-1999
			JP 2001525419 T	11-12-2001
EP 0394111	A	24-10-1990	FR 2645866 A1	19-10-1990
			AT 154035 T	15-06-1997
			CA 2014518 A1	17-10-1990
			DE 69030839 D1	10-07-1997
			DE 69030839 T2	20-11-1997
			DK 394111 T3	08-09-1997
			EP 0394111 A1	24-10-1990
			ES 2104593 T3	16-10-1997
			FR 2646161 A1	26-10-1990
			GR 3023691 T3	30-09-1997
			IL 94077 A	29-12-1994
			JP 2292246 A	03-12-1990
			JP 2716565 B2	18-02-1998
			US 5476962 A	19-12-1995
			US 5616745 A	01-04-1997
			US 5171678 A	15-12-1992

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)